

l3_tex_3

(TMXBKc42HG3HV4DHSkhJQj5HmxMH92Y2Fqm)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_tops_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_tops_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (((v1_tops_3 X1 X0) \wedge (v1_tops_1 X2 X0)) \Rightarrow (v1_tops_1 (k9_subset_1 \\ & (u1_struct_0 X0) X1 X2) X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\neg (v1_tops_1 X1 X0) \wedge (X1 = k1_xboole_0))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Rightarrow (r1_xboole_0 X1 X0) \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & X0)) \Rightarrow (m1_subset_1 (k9_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0. \forall X1. (r1_xboole_0 X0 X1) \Leftrightarrow (k3_xboole_0 X0 X1 = k1_xboole_0) \tag{6}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 \\ & X0))) \Rightarrow (\neg(((v1_tops_3 X1 X0) \wedge (v1_tops_1 X2 X0)) \vee ((v1_tops_1 X1 \\ & X0) \wedge (v1_tops_3 X2 X0))) \wedge (r1_xboole_0 X1 X2)))) \end{aligned}$$