

l3_uniroots

(TMN9esc8NiST4FHFJG9VzLE6EWCCSBBLfTw)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (k5_relat_1 (k5_relat_1 X2 X1) X0 = k5_relat_1 X2 X0)) \quad (1)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_tarski (k2_finseq_1 X0) (k2_finseq_1 X1)))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (5)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 \ k1_numbers \quad (7)$$

Assume the following.

$$m1_subset_1 \ k5_numbers \ (k1_zfmisc_1 \ k1_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(v6_membered \ X0) \Rightarrow (\forall X1.(m1_subset_1 \ X1 \ X0) \Rightarrow (v7_ordinal1 \ X1)) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v1_finseq_1 \ X0))) \Rightarrow \\ & \quad (\forall X1.(m2_subset_1 \ X1 \ k1_numbers \ k5_numbers) \Rightarrow (\forall X2. \\ & (m2_subset_1 \ X2 \ k1_numbers \ k5_numbers) \Rightarrow ((r1_xxreal_0 \ X2 \ X1) \Rightarrow \\ & (k5_relat_1 \ (k5_relat_1 \ X0 \ (k2_finseq_1 \ X1)) \ (k2_finseq_1 \ X2) = \\ & \quad k5_relat_1 \ X0 \ (k2_finseq_1 \ X2)))))) \end{aligned}$$