

l42_square_1

(TMGpw3wkV6DWFntNWxryGuf9dq1mVnHFemL)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $k6_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow ((k6_xcmplx_0 X0 X1 = k6_numbers) \Rightarrow (X0 = X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (v1_xcmplx_0 (k3_square_1 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xreal_0 (k6_square_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((X1 = k6_square_1 X0) \Leftrightarrow ((r1_xxreal_0 k6_numbers X1) \wedge (k3_square_1 X1 = X0))))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(r1_xxreal_0 k6_numbers X0) \wedge ((r1_xxreal_0 k6_numbers X1) \wedge ((X0 \neq X1) \wedge (k6_xcmplx_0 (k3_square_1 (k6_square_1 X0)) (k3_square_1 (k6_square_1 X1)) = k6_numbers)))))$$