

## l44\_fuzzy\_2

(TMPu1wFssQYVuTdsprSy4B4RNbq7se9PDPw)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k8\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_square\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k3\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_membered : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v2\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 k6\_numbers \\ & X2)) \Rightarrow (r1\_xxreal\_0 (k3\_xcmplx\_0 X0 X2) (k3\_xcmplx\_0 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 \\ & X1)) \Rightarrow (k8\_real\_1 X0 X1 = k3\_xcmplx\_0 X0 X1) \end{aligned} \quad (2)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (m1\_subset\_1 \\ & X1 k1\_numbers)) \Rightarrow (k1\_square\_1 X0 X1 = k3\_xxreal\_0 X0 X1) \end{aligned} \quad (4)$$

Assume the following.

$$v3\_membered k1\_numbers \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 \\ & X1)) \Rightarrow (m1\_subset\_1 (k8\_real\_1 X0 X1) k1\_numbers) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Rightarrow (k3\_xxreal\_0 X0 X1 = X0)) \wedge ((\neg r1\_xxreal\_0 X0 X1) \Rightarrow (k3\_xxreal\_0 X0 X1 = X1)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 X1)) \Rightarrow (k8\_real\_1 X0 X1 = k8\_real\_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (k3\_xxreal\_0 X0 X1 = k3\_xxreal\_0 X1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v3\_membered X0) \Rightarrow (v2\_membered X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v2\_membered X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (v1\_xxreal\_0 X1)) \quad (12)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 X1 k1\_numbers) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow ((r1\_xxreal\_0 k6\_numbers X2) \Rightarrow (k8\_real\_1 X2 (k1\_square\_1 X0 X1) = k1\_square\_1 (k8\_real\_1 X2 X0) (k8\_real\_1 X2 X1))))))$$