

l44_scmfsa10 (TMNwaYBnEZxn- HoF5Evnc5Ug4UiEttsGLza2)

October 27, 2020

Let $k5_scmfsa_2 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k4_scmfsa_2 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k5_scmfsa_2 X0 \neq k4_scmfsa_2 X1)) \quad (1)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (2)$$

Assume the following.

$$np_1 \in k4_ordinal1 \quad (3)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Leftrightarrow (X0 \in k4_ordinal1) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (v7_ordinal1 X0) \quad (6)$$

Theorem 1 $k5_scmfsa_2 k6_numbers \neq k4_scmfsa_2 np_1$.