

l45_fib_num4

(TMZdssz8JKoHyiBrJGTC4yKQZVFSAUfWTzd)

October 27, 2020

Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $np_5 : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xreal_0 X0 k6_numbers) \wedge ((\neg r1_xreal_0 \\ & X2 X1) \wedge (r1_xreal_0 (k3_xcmplx_0 X2 X0) (k3_xcmplx_0 X1 X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_5) \wedge (m2_subset_1 np_5 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_5 k5_numbers) \wedge (m1_subset_1 np_5 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (6)$$

Assume the following.

$$\neg r1_xreal_0 np_2 np_0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_boole_0 \quad (8)$$

Assume the following.

$$\neg r1_xreal_0 (k7_square_1 \ np_5) \ np_1 \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ k1_numbers) \Rightarrow (m1_subset_1 \ (k7_square_1 \ X0) \ k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 \ X0) \wedge (v1_xcmplx_0 \ X1)) \Rightarrow (k3_xcmplx_0 \ X0 \ X1 = k3_xcmplx_0 \ X1 \ X0) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ k1_numbers) \Rightarrow (v1_xreal_0 \ X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ k1_numbers) \Rightarrow (v1_xcmplx_0 \ X0) \quad (13)$$

Theorem 1

$$\neg r1_xreal_0 (k3_xcmplx_0 \ np_2 \ (k7_square_1 \ np_5)) \ (k3_xcmplx_0 \ np_2 \ np_1)$$