

# l45\_sin\_cos7

(TMdhprydFNSP1nuHecJSwHnrojY9P6HpDaE)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k12\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k11\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k9\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_square\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_0 : \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 k6\_numbers) \wedge ((\neg r1\_xxreal\_0 \\ & X2 X1) \wedge (r1\_xxreal\_0 (k7\_xcmplx\_0 X2 X0) (k7\_xcmplx\_0 X1 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 k6\_numbers) \wedge ((\neg r1\_xxreal\_0 \\ & X2 X1) \wedge (r1\_xxreal\_0 (k3\_xcmplx\_0 X2 X0) (k3\_xcmplx\_0 X1 X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k7\_xcmplx\_0 k6\_numbers X0 = k6\_numbers) \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2\_xreal\_0 \ np\_1) \wedge (m2\_subset\_1 \ np\_1 \ k1\_numbers \ k5\_numbers)) \wedge \\ & ((m1\_subset\_1 \ np\_1 \ k5\_numbers) \wedge (m1\_subset\_1 \ np\_1 \ k1\_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$v1\_xboole\_0 \ np\_0 \quad (7)$$

Assume the following.

$$k3\_xcmplx\_0 \ np\_0 \ np\_2 = np\_0 \quad (8)$$

Assume the following.

$$\neg r1\_xreal\_0 \ np\_2 \ np\_0 \quad (9)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xreal\_0 \ X0) \wedge (v1\_xreal\_0 \ X1)) \Rightarrow (k12\_binop\_2 \ X0 \ X1 = k7\_xcmplx\_0 \ X0 \ X1) \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xreal\_0 \ X0) \wedge (v1\_xreal\_0 \ X1)) \Rightarrow (k11\_binop\_2 \ X0 \ X1 = k3\_xcmplx\_0 \ X0 \ X1) \quad (12)$$

Assume the following.

$$\exists X0. (v1\_xboole\_0 \ X0) \wedge ((v1\_xcmplx\_0 \ X0) \wedge ((v1\_xreal\_0 \ X0) \wedge (v1\_xreal\_0 \ X0))) \quad (13)$$

Assume the following.

$$\forall X0. (v1\_xreal\_0 \ X0) \Rightarrow (\neg r1\_xreal\_0 \ (k9\_binop\_2 \ (k3\_square\_1 \ X0) \ np\_1) \ k6\_numbers) \quad (14)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xreal\_0 \ X0) \wedge (v1\_xreal\_0 \ X1)) \Rightarrow (v1\_xreal\_0 \ (k3\_xcmplx\_0 \ X0 \ X1)) \quad (15)$$

Assume the following.

$$\forall X0. (v1\_xreal\_0 \ X0) \Rightarrow (v1\_xreal\_0 \ (k3\_square\_1 \ X0)) \quad (16)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xreal\_0 \ X0) \wedge (v1\_xreal\_0 \ X1)) \Rightarrow (m1\_subset\_1 \ (k9\_binop\_2 \ X0 \ X1) \ k1\_numbers) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k9\_binop\_2 X0 X1 = k9\_binop\_2 X1 X0) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k11\_binop\_2 X0 X1 = k11\_binop\_2 X1 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(v1\_xcmplx\_0 X0) \quad (20)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (21)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(\neg(\neg r1\_xxreal\_0 X0 k6\_numbers)\wedge(r1\_xxreal\_0 (k12\_binop\_2 (k11\_binop\_2 np\_2 X0) (k9\_binop\_2 np\_1 (k3\_square\_1 X0))) k6\_numbers))$$