

l46_euclid_4

(TMct8jrCbv3XTPGLbg8uZ3Y9Tdayirn9RGq)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k8_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m2_finseq_2 X1 k1_numbers \\ & (k1_euclid X0)) \Rightarrow (\forall X2.(m2_finseq_2 X2 k1_numbers (k1_euclid \\ & X0)) \Rightarrow ((k12_euclid (k8_euclid X0 X1 X2) = k6_numbers) \Leftrightarrow (X1 = X2)))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m2_finseq_2 X1 k1_numbers \\ & (k1_euclid X0)) \Rightarrow (\forall X2.(m2_finseq_2 X2 k1_numbers (k1_euclid \\ & X0)) \Rightarrow (\neg(X1 \neq X2) \wedge (k12_euclid (k8_euclid X0 X1 X2) = k6_numbers)))) \end{aligned}$$