

l46_xreal_1 (TMGHteEdVGhax- uxGh4X8VRmZZqtQTrZnymx)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_xxreal_0 (k2_xcmplx_0 \\ & X0 X2) (k2_xcmplx_0 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow ((r1_xxreal_0 (\\ & k2_xcmplx_0 X0 X1) (k2_xcmplx_0 X2 X3)) \Leftrightarrow (r1_xxreal_0 (k6_xcmplx_0 \\ & X0 X2) (k6_xcmplx_0 X3 X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k6_xcmplx_0 X0 X0 = k6_numbers) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X1 X0) \wedge (r1_xxreal_0 (k6_xcmplx_0 X1 X0) k6_numbers)))$$