

# 148\_borsuk\_7 (TMVRuZjhZbVn- MdTE6p5ZQuDXLrWfHwRuMfd)

October 27, 2020

Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_pre\_topc : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_topmetr : \iota$  be given. Let  $k5\_toprealb : \iota \Rightarrow \iota$  be given. Let  $c15\_borsuk\_7 : \iota$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(l1\_pre\_topc X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (u1\_struct\_0 (k1\_pre\_topc X0 X1) = X1)) \quad (1)$$

Assume the following.

$$u1\_struct\_0 k3\_topmetr = k1\_numbers \quad (2)$$

Assume the following.

$$(v2\_pre\_topc k3\_topmetr) \wedge (l1\_pre\_topc k3\_topmetr) \quad (3)$$

Assume the following.

$$m1\_subset\_1 c15\_borsuk\_7 (k1\_zfmisc\_1 k1\_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 k1\_numbers)) \Rightarrow (k5\_toprealb X0 = X0) \quad (5)$$

## Theorem 1

$$u1\_struct\_0 (k1\_pre\_topc k3\_topmetr (k5\_toprealb c15\_borsuk\_7)) = c15\_borsuk\_7$$