

l48_subset_1 (TMVmKCypq- buoVpdwA2CGXuHs7ZZd8jmtKXP)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \neg v1_xboole_0 (k1_zfmisc_1 X0) \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_zfmisc_1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (r1_tarski X2 X0)) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (X1 \in X0))) \wedge ((v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (v1_xboole_0 X1))) \tag{4}$$

Theorem 1

$$\forall X0. \forall X1. (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \Rightarrow (m1_subset_1 X0 (k1_zfmisc_1 X1))$$