

l49_modal_1 (TMVN- BXA3EiTbbKrYrifhGEYPyLG2amcTYoN)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_trees_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k8_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$k2_trees_1 \ np_1 = k2_tarski \ k1_xboole_0 \ (k12_finseq_1 \ k5_numbers \ k6_numbers) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 \ (k2_funcop_1 \ X0 \ X1) = X0) \wedge (r1_tarski \ (k10_xtuple_0 \ (k2_funcop_1 \ X0 \ X1)) \ (k1_tarski \ X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 \ X0) \wedge (m1_subset_1 \ X2 \ X0)) \Rightarrow (k8_funcop_1 \ X0 \ X1 \ X2 = k2_funcop_1 \ X1 \ X2) \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\neg v1_finset_1 k4_ordinal1 \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_finset_1 X0)\wedge(\neg v1_xboole_0 X1))\Rightarrow(\neg v1_finset_1 (k2_zfmisc_1 X1 X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X2 X0))\Rightarrow((v1_funct_1 (k8_funcop_1 X0 X1 X2))\wedge((v1_funct_2 (k8_funcop_1 X0 X1 X2) X1 X0)\wedge(m1_subset_1 (k8_funcop_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X1 X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge((m1_subset_1 X2 X0)\wedge(m1_subset_1 X3 X1))))\Rightarrow(m1_subset_1 (k1_domain_1 X0 X1 X2 X3) (k2_zfmisc_1 X0 X1)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k2_tarski X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 = X0)\vee(X3 = X1))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (14)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v1_finset_1 X0) \quad (15)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k5_numbers)\Rightarrow(\forall X1.(m1_subset_1 X1 k5_numbers)\Rightarrow(k12_finseq_1 k5_numbers k6_numbers \in k1_relset_1 (k2_trees_1 np_1) (k8_funcop_1 (k2_zfmisc_1 k5_numbers k5_numbers) (k2_trees_1 np_1) (k1_domain_1 k5_numbers k5_numbers X0 X1))))$$