

l4_rlvect_5 (TMbPgKsQ- pANaYrp89D8QAXaU3ZBctr5hddy)

October 27, 2020

Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (k2_xboole_0 (k4_xboole_0 X1 (k1_tarski X0)) (k1_tarski X0) = X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. (X1 \in X0) \Rightarrow (k2_xboole_0 (k6_subset_1 X0 (k1_tarski X1)) (k1_tarski X1) = X0)$$