

l52_xxreal_3

(TMYaJgbChpwJruh25X9g2ZCKRSFg2JVK47f)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k1_xxreal_3 X0 k6_numbers = X0) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 (k1_xxreal_3 \\ & X0 X2) (k1_xxreal_3 X1 X2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v1_xxreal_0 (k1_xxreal_3 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow(k1_xxreal_3 X0 X1 = k1_xxreal_3 X1 X0) \quad (8)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0)\Rightarrow(\forall X1.(v1_xxreal_0 X1)\Rightarrow(\neg(r1_xxreal_0 X0 k6_numbers)\wedge(\neg r1_xxreal_0 k6_numbers X1)\wedge(r1_xxreal_0 k6_numbers (k1_xxreal_3 X0 X1))))$$