

# l53\_toprealb (TMVJeWN- SnuSyvsYxxTVjZ1ZRdZDSw14nfbJ)

October 27, 2020

Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k11\_toprealb : \iota \Rightarrow \iota$  be given. Let  $k9\_toprealb : \iota$  be given. Let  $k2\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $np\_2 : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $m1\_pre\_topc : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $l1\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $k8\_toprealb : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $v5\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $v1\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 \ np\_2) \wedge (m2\_subset\_1 \ np\_2 \ k1\_numbers \ k5\_numbers)) \wedge \\ & ((m1\_subset\_1 \ np\_2 \ k5\_numbers) \wedge (m1\_subset\_1 \ np\_2 \ k1\_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (2)$$

Assume the following.

$$\forall X0. (l1\_pre\_topc \ X0) \Rightarrow (\forall X1. (m1\_pre\_topc \ X1 \ X0) \Rightarrow (l1\_pre\_topc \ X1)) \quad (3)$$

Assume the following.

$$\forall X0. (l1\_rltopsp1 \ X0) \Rightarrow ((l1\_rlvect\_1 \ X0) \wedge (l1\_pre\_topc \ X0)) \quad (4)$$

Assume the following.

$$\forall X0. (l1\_pre\_topc \ X0) \Rightarrow (l1\_struct\_0 \ X0) \quad (5)$$

Assume the following.

$$m1\_subset\_1 \ k9\_toprealb \ (u1\_struct\_0 \ (k8\_toprealb \ np\_2)) \quad (6)$$

Assume the following.

$$\forall X0.(v7\_ordinal1\ X0)\Rightarrow(m1\_pre\_topc\ (k8\_toprealb\ X0)\ (k15\_euclid\ X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v7\_ordinal1\ X0)\Rightarrow((v5\_rltopsp1\ (k15\_euclid\ X0))\wedge\ (l1\_rltopsp1\ (k15\_euclid\ X0))) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1\ X0\ (u1\_struct\_0\ (k8\_toprealb\ np\_2)))\Rightarrow\ ((v1\_pre\_topc\ (k11\_toprealb\ X0))\wedge(m1\_pre\_topc\ (k11\_toprealb\ X0)\ (k8\_toprealb\ np\_2))) \quad (9)$$

Assume the following.

$$\forall X0.(l1\_struct\_0\ X0)\Rightarrow(k2\_struct\_0\ X0 = u1\_struct\_0\ X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1\ X0\ k4\_ordinal1)\Rightarrow(v7\_ordinal1\ X0) \quad (11)$$

**Theorem 1**

$$u1\_struct\_0\ (k11\_toprealb\ k9\_toprealb) = k2\_struct\_0\ (k11\_toprealb\ k9\_toprealb)$$