

l56_group_1 (TMd- wqcjg7RZYGidvBCEwUDcXj7YzoTNnWWG)

October 27, 2020

Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k19_binop_2 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_int_1 : \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k5_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $c5_xreal_0 : \iota$ be given. Let $k1_arytm_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $c3_xreal_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((k1_int_1 X0 = X0) \Leftrightarrow (v1_int_1 X0)) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k2_xcmplx_0 X0 X1) X2) \Leftrightarrow (r1_xxreal_0 \\ X0 (k6_xcmplx_0 X2 X1)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (5)$$

Assume the following.

$$k2_xcmplx_0 (k4_xcmplx_0 np_1) np_1 = np_0 \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(m1_subset_1 X1 k1_numbers))\Rightarrow (k5_real_1 X0 X1 = k6_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.(v1_int_1 X0)\Rightarrow(k19_binop_2 X0 = k4_xcmplx_0 X0) \quad (10)$$

Assume the following.

$$(c5_xreal_0 = k4_xcmplx_0 np_1)\wedge(k1_arytm_0 c3_xreal_0 c5_xreal_0 = k6_numbers) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (12)$$

Assume the following.

$$\forall X0.(v1_int_1 X0)\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(v1_int_1 (k4_xcmplx_0 X0))) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_int_1 X1)\Rightarrow((X1 = k1_int_1 X0)\Leftrightarrow((r1_xxreal_0 X1 X0)\wedge(\neg r1_xxreal_0 X1 (k5_real_1 X0 np_1)))) \quad (14)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (15)$$

Assume the following.

$$\forall X0.(v1_int_1 X0)\Rightarrow(v1_xreal_0 X0) \quad (16)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_int_1 X0) \quad (17)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (18)$$

Theorem 1

$$\forall X0.(v1_int_1 X0)\Rightarrow(\neg(\neg r1_xxreal_0 k6_numbers X0)\wedge((X0\neq k19_binop_2 np_1)\wedge(r1_xxreal_0 (k19_binop_2 np_1) X0)))$$