

l58_orders_1

(TMZoym5T1vrkXVkAGyNM4kow3Fe1t118hyZ)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_relat_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_relat_1 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (v1_relat_1 (k2_relat_1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(v1_relat_1 X1) \Rightarrow ((X1 = k2_relat_1 X0) \Leftrightarrow (\forall X2.\forall X3.(k4_tarski X2 X3 \in X1) \Leftrightarrow (k4_tarski X3 X2 \in X0)))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(r1_relat_2 X0 X1) \Leftrightarrow (\forall X2.(X2 \in X1) \Rightarrow (k4_tarski X2 X2 \in X0))) \quad (3)$$

Theorem 1

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(r1_relat_2 X0 X1) \Rightarrow (r1_relat_2 (k2_relat_1 X0) X1))$$