

l5_catalan2

(TMTY88bvYPbsJbsxu8g8Foqb7vACanAJDKy)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_ordinal1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_ordinal1 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k5_relat_1 (k2_funcop_1 X0 X1) X2 = k2_funcop_1 (k3_xboole_0 X0 X2) X1 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v3_ordinal1 X0) \wedge (v3_ordinal1 X1)) \Rightarrow ((r1_ordinal1 X0 X1) \Leftrightarrow (r1_tarski X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k5_relset_1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_funct_1 (k7_funcop_1 X0 X1))\wedge((v1_funct_2 (k7_funcop_1 X0 X1) X0 (k1_tarSKI X1))\wedge(m1_subset_1 (k7_funcop_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarSKI X1)))))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (8)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v3_ordinal1 X0) \quad (9)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(\forall X2.(v7_ordinal1 X2)\Rightarrow((r1_xreal_0 X0 X1)\Rightarrow(k5_relset_1 X1 (k1_tarSKI X2) (k7_funcop_1 X1 X2) X0 = k7_funcop_1 X0 X2))))$$