

l5_finseq_7 (TMP rMZY-
dGq7TQAo5Py2XE538WTUrnv3q3rP)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k7_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7_ordinal1 X2) \Rightarrow (k7_nat_d (k7_nat_d X0 X1) X2 = k7_nat_d X0 (k2_xcmplx_0 \\ & X1 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \tag{2}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \tag{4}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \tag{5}$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xcmplx_0 X0) \tag{6}$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k7_nat_d (k7_nat_d X0 X1) np_1 = k7_nat_d (k7_nat_d X0 np_1) X1))$$