

l5_fomodel0

(TMHu9KSpZKyehtT1jpfwxoXNyrW6AWUiS67)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((m1_finseq_1 X1 X0) \Leftrightarrow (r1_tarski (k10_xtuple_0 X1) X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((v5_relat_1 X1 X0) \Leftrightarrow (r1_tarski (k10_xtuple_0 X1) X0)) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((v5_relat_1 X1 X0) \wedge (v1_finseq_1 X1)) \Rightarrow (m2_finseq_1 X1 X0))$$