

# l5\_wsierp\_1 (TMY- hYWi8LwwPXyYPTcRtawaDnyzjKe3kRnS)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 \\ & X1 X2)) \Rightarrow (r1\_xxreal\_0 X1 (k2\_xcmplx\_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 (k2\_xcmplx\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 \\ & (k6\_xcmplx\_0 X0 X1) X2)))) \end{aligned} \quad (2)$$

## Theorem 1

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 \\ & X2 X1)) \Rightarrow ((r1\_xxreal\_0 X2 (k2\_xcmplx\_0 X0 X1)) \wedge (r1\_xxreal\_0 (k6\_xcmplx\_0 \\ & X2 X0) X1)))) \end{aligned}$$