

l63_series_3

(TMQsR8Gzg4RHMMomJX5JPoXVqfnjTBBSxX3a)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_square_1 : \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_series_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (r1_xxreal_0 k6_numbers (k3_xcmplx_0 X0 X0)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow (r1_xxreal_0 k6_numbers \\ & (k1_seq_1 X0 X1))) \Rightarrow (\forall X1.(m1_subset_1 X1 k5_numbers) \Rightarrow (\\ & r1_xxreal_0 k6_numbers (k1_seq_1 (k3_series_1 X0) X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k5_square_1 X0 = k3_square_1 X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k7_real_1 X0 X1) k1_numbers) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (m1_subset_1 (k1_seq_1 X0 X1) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_square_1 X0 = k3_xcmplx_0 X0 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v5_relat_1 X0 k1_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.((v1_funct_1 \\ & X1) \wedge ((v1_funct_2 X1 k5_numbers k1_numbers) \wedge (m1_subset_1 X1 (\\ & k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow (\forall X2. \\ & ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k5_numbers k1_numbers) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow (\forall X3. \\ & ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 k5_numbers k1_numbers) \wedge (m1_subset_1 \\ & X3 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((\forall X4. \\ & (m1_subset_1 X4 k5_numbers) \Rightarrow (k1_seq_1 X1 X4 = k5_square_1 (k7_real_1 \\ & (k1_seq_1 X2 X4) (k1_seq_1 X3 X4)))) \Rightarrow (r1_xxreal_0 k6_numbers (\\ & k1_seq_1 (k3_series_1 X1) X0)))))) \end{aligned}$$