

## l64\_cqc\_the3

(TMPV4PZtcoYEhFgodz4ZxqaowwWBiz1uadv)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $r7\_cqc\_the3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_cqc\_the1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow ((v2\_cqc\_the1 (k8\_cqc\_lang X0 (k6\_cqc\_lang \\ X0 X1) (k6\_cqc\_lang X0 X2)) X0) \Leftrightarrow (v2\_cqc\_the1 (k8\_cqc\_lang X0 X2 \\ X1) X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow ((r7\_cqc\_the3 X0 X1 X2) \Leftrightarrow ((v2\_cqc\_the1 ( \\ k8\_cqc\_lang X0 X1 X2) X0) \wedge (v2\_cqc\_the1 (k8\_cqc\_lang X0 X2 X1) X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\neg v1\_xboole\_0 (k3\_cqc\_lang X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((m1\_qc\_lang1 X0) \wedge (m1\_subset\_1 X1 (k3\_cqc\_lang \\ X0))) \Rightarrow (m2\_subset\_1 (k6\_cqc\_lang X0 X1) (k9\_qc\_lang1 X0) (k3\_cqc\_lang \\ X0)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (m1\_subset\_1 (k3\_cqc\_lang X0) (k1\_zfmisc\_1 (k9\_qc\_lang1 X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (v1\_xboole\_0 X1)) \quad (7)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow ((r7\_cqc\_the3 X0 X1 X2) \Rightarrow (r7\_cqc\_the3 X0 \\ (k6\_cqc\_lang X0 X1) (k6\_cqc\_lang X0 X2)))))) \end{aligned}$$