

l64_graphsp

(TMR9K2NU2wZS5HrzyusTfSXQsuYdioCexCV)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xreal_0 X0 (k2_xcmplx_0 X1 X2)) \Leftrightarrow (r1_xreal_0 \\ & (k6_xcmplx_0 X0 X1) X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (r1_xreal_0 X0 (k2_xcmplx_0 X0 X1))) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k9_real_1 X0 X1 = k6_xcmplx_0 X0 X1) \tag{4}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \tag{6}$$

Assume the following.

$$(\neg v1_xboole_0\ k4_ordinal1) \wedge (v3_ordinal1\ k4_ordinal1) \quad (7)$$

Assume the following.

$$\neg v1_xboole_0\ k1_numbers \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0\ X0) \wedge ((\neg v1_xboole_0\ X1) \wedge \\ (m1_subset_1\ X1\ (k1_zfmisc_1\ X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ X2\ X0\ X1) \Rightarrow (m1_subset_1\ X2\ X0)) \end{aligned} \quad (9)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1\ X0\ k5_numbers) \wedge (v7_ordinal1\ X1)) \Rightarrow (k2_nat_1\ X0\ X1 = k2_nat_1\ X1\ X0) \quad (11)$$

Assume the following.

$$\forall X0. (m1_subset_1\ X0\ k4_ordinal1) \Rightarrow (v7_ordinal1\ X0) \quad (12)$$

Assume the following.

$$\forall X0. (m1_subset_1\ X0\ k1_numbers) \Rightarrow (v1_xreal_0\ X0) \quad (13)$$

Theorem 1

$$\begin{aligned} \forall X0. (m2_subset_1\ X0\ k1_numbers\ k5_numbers) \Rightarrow (\forall X1. \\ (m2_subset_1\ X1\ k1_numbers\ k5_numbers) \Rightarrow (r1_xreal_0\ (k9_real_1 \\ X0\ X1)\ X0)) \end{aligned}$$