

l64_xcmlx_1 (TMLKn- FzhkVYYvf1onYWchTEDmgkaqqvmnxb)

October 27, 2020

Let $v1_xcmlx_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcmlx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmlx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmlx_0 X0) \Rightarrow (\forall X1.(v1_xcmlx_0 X1) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (X1 = k7_xcmlx_0 X1 (k7_xcmlx_0 X0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmlx_0 X0) \Rightarrow (\forall X1.(v1_xcmlx_0 X1) \Rightarrow (\forall X2.(v1_xcmlx_0 X2) \Rightarrow (k7_xcmlx_0 X0 (k7_xcmlx_0 X1 X2) = k7_xcmlx_0 (k3_xcmlx_0 X0 X2) X1))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmlx_0 X0) \wedge (v1_xcmlx_0 X1)) \Rightarrow (v1_xcmlx_0 (k3_xcmlx_0 X0 X1)) \quad (3)$$

Theorem 1

$$\forall X0.(v1_xcmlx_0 X0) \Rightarrow (\forall X1.(v1_xcmlx_0 X1) \Rightarrow (\forall X2.(v1_xcmlx_0 X2) \Rightarrow ((X0 \neq k6_numbers) \Rightarrow (k7_xcmlx_0 X1 X2 = k7_xcmlx_0 (k3_xcmlx_0 X1 X0) (k3_xcmlx_0 X2 X0))))))$$