

l65_cqc_the3
(TMMU72hrKaFc3mUUaVnVFqhWQeHuAGd5Dxj)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $r7_cqc_the3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_cqc_the1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow ((v2_cqc_the1\ (k8_cqc_lang\ X0\ (k6_cqc_lang \\ X0\ X1)\ (k6_cqc_lang\ X0\ X2))\ X0) \Leftrightarrow (v2_cqc_the1\ (k8_cqc_lang\ X0\ X2 \\ X1)\ X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow ((r7_cqc_the3\ X0\ X1\ X2) \Leftrightarrow ((v2_cqc_the1\ (\\ k8_cqc_lang\ X0\ X1\ X2)\ X0) \wedge (v2_cqc_the1\ (k8_cqc_lang\ X0\ X2\ X1)\ X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0\ X0) \wedge ((\neg v1_xboole_0\ X1) \wedge \\ (m1_subset_1\ X1\ (k1_zfmisc_1\ X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2\ X0\ X1) \Leftrightarrow (m1_subset_1\ X2\ X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0) \Rightarrow (\neg v1_xboole_0\ (k3_cqc_lang\ X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((m1_qc_lang1\ X0) \wedge (m1_subset_1\ X1\ (k3_cqc_lang \\ X0))) \Rightarrow (m2_subset_1\ (k6_cqc_lang\ X0\ X1)\ (k9_qc_lang1\ X0)\ (k3_cqc_lang \\ X0)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k3_cqc_lang X0) (k1_zfmisc_1 (k9_qc_lang1 X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ & X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ & X0) (k3_cqc_lang X0)) \Rightarrow ((r7_cqc_the3 X0 (k6_cqc_lang X0 X1) (k6_cqc_lang \\ & X0 X2)) \Rightarrow (r7_cqc_the3 X0 X1 X2)))) \end{aligned}$$