

l65_fib_num4
(TMPnJgKG8xZFoUzv9ffcca6Wn6ffmhBsNdR)

October 27, 2020

Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $np_5 : \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $c1_xreal_1 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xreal_0 X0 X1) \Leftrightarrow (r1_xreal_0 (k6_xcmplx_0 \\ & X0 X2) (k6_xcmplx_0 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_5) \wedge (m2_subset_1 np_5 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_5 k5_numbers) \wedge (m1_subset_1 np_5 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\neg r1_xreal_0 (k7_square_1 np_5) np_1 \quad (3)$$

Assume the following.

$$m1_subset_1 np_1 k1_numbers \quad (4)$$

Assume the following.

$$\neg r1_xreal_0 (k7_square_1 np_5) k6_numbers \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k7_square_1 X0) k1_numbers) \quad (6)$$

Assume the following.

$$c1_xreal_1 = k6_numbers \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$(-r1_xreal_0 (k6_xcmplx_0 (k7_square_1 np_5) np_1) (k6_xcmplx_0 np_1 np_1))^{(-r1_xreal_0 (k7_square_1 np_5) k6_numbers)}$$