

l65_sin_cos (TMEx- aBTEpgB3PmN2XAjxqk2CLuCy5tzE4VE)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0 : \iota \Rightarrow \iota. \exists X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 \\ & X1 k5_numbers k1_numbers) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers k1_numbers)))) \wedge (\forall X2. (m2_subset_1 X2 k1_numbers \\ & k5_numbers) \Rightarrow (k1_seq_1 X1 X2 = X0 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1_funct_1 X1) \wedge ((v1_funct_2 \\ & X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))) \wedge (v7_ordinal1 X2)) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2)) \end{aligned} \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 \\ & X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \end{aligned} \quad (5)$$

Assume the following.

$$(\neg v1_xboole_0\ k4_ordinal1) \wedge (v3_ordinal1\ k4_ordinal1) \quad (6)$$

Assume the following.

$$v3_membered\ k1_numbers \quad (7)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1) \Rightarrow (v7_ordinal1\ X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0)) \Rightarrow (v1_xboole_0\ X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))) \Rightarrow (v1_relat_1\ X2) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered\ X1) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))) \Rightarrow (v3_valued_0\ X2)) \quad (12)$$

Theorem 1

$$\begin{aligned} & \exists X0.((v1_funct_1\ X0) \wedge ((v1_funct_2\ X0\ k5_numbers\ k1_numbers) \wedge \\ & (m1_subset_1\ X0\ (k1_zfmisc_1\ (k2_zfmisc_1\ k5_numbers\ k1_numbers)))))) \wedge \\ & (\forall X1.(m2_subset_1\ X1\ k1_numbers\ k5_numbers) \Rightarrow (k8_nat_1\ k1_numbers\ X0\ X1 = k7_real_1\ (k8_real_1\ np_2\ X1)\ np_1)) \end{aligned}$$