

l67_xreal_1

(TMRgmVhDr4UfDDV5tayBRsXng7tsvZHzR1s)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 X1) \wedge ((\neg v3_xxreal_0 X1) \wedge (\neg v2_xxreal_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v2_xxreal_0 X0)) \Rightarrow (v2_xxreal_0 X1))) \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 X2 X1) \wedge (r1_xxreal_0 (k3_xcmplx_0 X2 X0) (k3_xcmplx_0 X1 X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k5_xcmplx_0 (k5_xcmplx_0 X0) = X0) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k5_xcmplx_0 X0)) \wedge (v1_xreal_0 (k5_xcmplx_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.((\neg v2_xxreal_0 X0) \wedge (v1_xreal_0 X0)) \Rightarrow ((v1_xcmplx_0 (k5_xcmplx_0 X0)) \wedge (\neg v2_xxreal_0 (k5_xcmplx_0 X0))) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k7_xcmplx_0 X0 X1 = k3_xcmplx_0 X0 (k5_xcmplx_0 X1))) \quad (10)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v3_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v2_xxreal_0 X0))) \quad (11)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v2_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X0))) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (13)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((\neg r1_xxreal_0 X2 X1) \wedge (r1_xxreal_0 (k7_xcmplx_0 X2 X0) (k7_xcmplx_0 X1 X0)))))))$$