

l69_modal_1

(TMS6G8Kag5mBnCxBiYtQUBdxXdzRNAWSESw)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k4_modal_1 : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_mcart_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k8_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (1)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((m1_subset_1 X2 \\ & (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X3 (k1_zfmisc_1 X1))) \Rightarrow (k8_mcart_1 \\ & X0 X1 X2 X3 = k2_zfmisc_1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1_xboole_0 X0) \wedge \\ & ((m1_subset_1 X1 X0) \wedge ((m1_subset_1 X2 X0) \wedge (m1_subset_1 X3 X0)))) \Rightarrow \\ & (k8_domain_1 X0 X1 X2 X3 = k1_enumset1 X1 X2 X3) \end{aligned} \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ & ((\neg v1_xboole_0 X1) \wedge ((m1_subset_1 X2 X0) \wedge (m1_subset_1 X3 X1)))) \Rightarrow \\ & (k1_domain_1 X0 X1 X2 X3 = k4_tarSKI X2 X3) \end{aligned} \quad (9)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\neg v1_xboole_0 (k1_enumset1 X0 X1 X2) \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ & ((m1_subset_1 X1 X0) \wedge ((m1_subset_1 X2 X0) \wedge (m1_subset_1 X3 X0)))) \Rightarrow \\ & (m1_subset_1 (k8_domain_1 X0 X1 X2 X3) (k1_zfmisc_1 X0)) \end{aligned} \quad (12)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ & ((\neg v1_xboole_0 X1) \wedge ((m1_subset_1 X2 X0) \wedge (m1_subset_1 X3 X1)))) \Rightarrow \\ & (m1_subset_1 (k1_domain_1 X0 X1 X2 X3) (k2_zfmisc_1 X0 X1)) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} k4_modal_1 = k8_mcart_1 k5_numbers k1_numbers (k8_domain_1 k5_numbers \\ k6_numbers np_1 np_2) k5_numbers \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(X3 = k1_enumset1 \\ & X0 X1 X2) \Leftrightarrow (\forall X4.(X4 \in X3) \Leftrightarrow (\neg(X4 \neq X0) \wedge ((X4 \neq X1) \wedge (X4 \neq X2)))) \end{aligned} \quad (16)$$

Theorem 1

$$\begin{aligned} & m1_subset_1 (k1_domain_1 k5_numbers k5_numbers k6_numbers k6_numbers) \\ & k4_modal_1 \end{aligned}$$