

l6_lukasi_1 (TM- NXB9gF84kzTFH34RxyNE2zNBP9LkXqsZe)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $k8_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_cqc_the1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1\ X0) \\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1\ X0) \\ (k3_cqc_lang\ X0)) \Rightarrow (((X1 \in k4_cqc_the1\ X0) \wedge (k8_cqc_lang\ X0\ X1\ X2 \in k4_cqc_the1\ X0)) \Rightarrow (X2 \in k4_cqc_the1\ X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0\ X0) \wedge ((\neg v1_xboole_0\ X1) \wedge \\ (m1_subset_1\ X1\ (k1_zfmisc_1\ X0)))) \Rightarrow (\forall X2.(m2_subset_1\ X2\ X0\ X1) \Leftrightarrow (m1_subset_1\ X2\ X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1\ X0) \\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1\ X0) \\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X3.(m2_subset_1\ X3\ (k9_qc_lang1\ X0) \\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X4.(m2_subset_1\ X4\ (k9_qc_lang1\ X0) \\ (k3_cqc_lang\ X0)) \Rightarrow (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ X1\ X2) \\ (k8_cqc_lang\ X0\ X3\ X2))\ X4)\ (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ X3\ X1)\ X4) \in k4_cqc_the1\ X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0) \Rightarrow (\neg v1_xboole_0\ (k3_cqc_lang\ X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1\ X0) \wedge ((m1_subset_1\ X1\ (k3_cqc_lang\ X0)) \wedge (m1_subset_1\ X2\ (k3_cqc_lang\ X0)))) \Rightarrow (m2_subset_1\ (k8_cqc_lang\ X0\ X1\ X2)\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0) \Rightarrow (m1_subset_1\ (k3_cqc_lang\ X0)\ (k1_zfmisc_1\ (k9_qc_lang1\ X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0)) \Rightarrow (v1_xboole_0\ X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X3.(m2_subset_1\ X3\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X4.(m2_subset_1\ X4\ (k9_qc_lang1\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ X1\ (k8_cqc_lang\ X0\ X2\ X3))\ (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ X4\ X2)\ (k8_cqc_lang\ X0\ X1\ (k8_cqc_lang\ X0\ X4\ X3))) \in k4_cqc_the1\ X0)))))) \end{aligned}$$