

## l6\_orders\_1

(TMYRoUusk6iNB6YeZEGS1R6gs46ikDz9KMe)

October 27, 2020

Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Rightarrow (k2\_xboole\_0 X0 X1 = X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow (k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow (m1\_subset\_1 (k2\_relset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \quad (5)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Rightarrow (k1\_relat\_1 X0 = k2\_xboole\_0 (k9\_xtuple\_0 X0) (k10\_xtuple\_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow( (v1\_partfun1 X1 X0)\Leftrightarrow(k1\_relset\_1 X0 X1 = X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k2\_xboole\_0 X0 X1 = k2\_xboole\_0 X1 X0 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (10)$$

**Theorem 1**

$$\forall X0.\forall X1.((v1\_partfun1 X1 X0)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0))))\Rightarrow(k1\_relat\_1 X1 = X0)$$