

l6_taylor_2

(TMKopeSa6hu2BWKLPjPwuLfqF1tdkSJvpQo)

October 27, 2020

Let $v3_rcomp_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow (k2_partfun1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (5)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 \ k1_numbers \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 \ X2 \ (k1_zfmisc_1 \\ & (k2_zfmisc_1 \ X0 \ X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 \ X2 \ X0 \\ & X1) \Leftrightarrow (X0 = k1_relset_1 \ X0 \ X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 \\ & X2 \ X0 \ X1) \Leftrightarrow (X2 = k1_xboole_0)))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 \ X0 \ X1 = k3_xboole_0 \ X1 \ X0 \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 \ X2 \ (k1_zfmisc_1 \\ & (k2_zfmisc_1 \ X0 \ X1))) \Rightarrow ((v4_relat_1 \ X2 \ X0) \wedge (v5_relat_1 \ X2 \ X1)) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 \ X2 \ (k1_zfmisc_1 \\ & (k2_zfmisc_1 \ X0 \ X1))) \Rightarrow (v1_relat_1 \ X2) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v3_rcomp_1 \ X0) \wedge (m1_subset_1 \ X0 \ (k1_zfmisc_1 \ k1_numbers))) \Rightarrow \\ & (\forall X1.((v1_funct_1 \ X1) \wedge ((v1_funct_2 \ X1 \ k1_numbers \ k1_numbers) \wedge \\ & (m1_subset_1 \ X1 \ (k1_zfmisc_1 \ (k2_zfmisc_1 \ k1_numbers \ k1_numbers)))))) \Rightarrow \\ & (k9_xtuple_0 \ (k2_partfun1 \ k1_numbers \ k1_numbers \ X1 \ X0) = X0) \end{aligned}$$