

l6_topreal2
(TMcSfCK3earkY4o2D1atBwCNg621D6fHhUy)

October 27, 2020

Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $c1_axioms : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \quad (1)$$

Assume the following.

$$m1_subset_1 np_1 k1_numbers \quad (2)$$

Assume the following.

$$m1_subset_1 k6_numbers k1_numbers \quad (3)$$

Assume the following.

$$c1_axioms = k6_numbers \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (5)$$

Theorem 1 $k17_euclid (k19_euclid k6_numbers np_1) = k6_numbers$.