

l6_toprealb (TMXcuGHuFfQdbd- VEj6ezMbEyqYC5iMmMqhM)

October 27, 2020

Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $np_2 : \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 \ np_3) \wedge (m2_subset_1 \ np_3 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_3 \ k5_numbers) \wedge (m1_subset_1 \ np_3 \ k1_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 \ np_2) \wedge (m2_subset_1 \ np_2 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_2 \ k5_numbers) \wedge (m1_subset_1 \ np_2 \ k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 \ np_1) \wedge (m2_subset_1 \ np_1 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_1 \ k5_numbers) \wedge (m1_subset_1 \ np_1 \ k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k6_xcmplx_0 \ (k7_xcmplx_0 \ np_3 \ np_2) \ (k7_xcmplx_0 \ np_1 \ np_2) = np_1 \quad (4)$$

Assume the following.

$$k6_xcmplx_0 \ np_2 \ (k7_xcmplx_0 \ np_1 \ np_2) = k7_xcmplx_0 \ np_3 \ np_2 \quad (5)$$

Assume the following.

$$r1_xxreal_0 \ np_1 \ np_1 \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 \ X0 \ k1_numbers) \wedge (v1_xreal_0 \ X1)) \Rightarrow (k9_real_1 \ X0 \ X1 = k6_xcmplx_0 \ X0 \ X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(k10_real_1 X0 X1 = k7_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k7_xcmplx_0 X0 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers)\wedge(v1_xreal_0 X1))\Rightarrow(m1_subset_1 (k9_real_1 X0 X1) k1_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$r1_xxreal_0 (k9_real_1 (k10_real_1 np_3 np_2) (k10_real_1 np_1 np_2)) np_1$$