

## l6\_trees\_4

(TMME3i4V9z2K5JqQySE8cskFPTFmg2MQFmQ)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1\_relat\_1 X2) \wedge ((v5\_relat\_1 X2 X0) \wedge (v1\_funct\_1 X2))) \Rightarrow ((X1 \in k9\_xtuple\_0 X2) \Rightarrow (k1\_funct\_1 X2 X1 \in X0)) \quad (1)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((\neg r1\_xxreal\_0 (k3\_finseq\_1 X1) X0) \Rightarrow ((k2\_nat\_1 X0 np\_1 \in k4\_finseq\_1 X1) \wedge (k1\_funct\_1 X1 (k2\_nat\_1 X0 np\_1) \in k10\_xtuple\_0 X1)))))) \quad (6)$$

Assume the following.

$$v6\_membered k4\_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0) \Rightarrow ((v1\_funct\_1 X1) \wedge ((v1\_finseq\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 X1)) \Rightarrow (k2\_nat\_1 X0 X1 = k2\_nat\_1 X1 X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow (v5\_relat\_1 X1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v6\_membered X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (v7\_ordinal1 X1)) \quad (12)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers) \Rightarrow (\forall X1.\forall X2.(m2\_finseq\_1 X2 X1) \Rightarrow ((\neg r1\_xxreal\_0 (k3\_finseq\_1 X2) X0) \Rightarrow (k1\_funct\_1 X2 (k2\_nat\_1 X0 np\_1) \in X1)))$$