

l75_fib_num4

(TMJ6KDuJM4Pihu6EzShrRHX7DCMAb1RJMcD)

October 27, 2020

Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_6 : \iota$ be given. Let $np_10 : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $np_5 : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k3_xcmplx_0 X0 (k7_xcmplx_0 np_1 X1) = k7_xcmplx_0 X0 X1)) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_5) \wedge (m2_subset_1 np_5 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_5 k5_numbers) \wedge (m1_subset_1 np_5 k1_numbers)) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k13_complex1 X0 X1 = k7_xcmplx_0 X0 X1) \quad (4)$$

Assume the following.

$$\neg r1_xxreal_0 (k13_complex1 np_6 np_10) (k3_xcmplx_0 (k6_xcmplx_0 (k7_square_1 np_5) np_1) (k13_complex1 np_1 (k7_square_1 np_5))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (v1_xcmplx_0 (k6_xcmplx_0 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k7_square_1 X0) k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xcmplx_0 X0) \quad (8)$$

Theorem 1

$$\neg r1_xreal_0 (k13_complex1 np_6 np_10) (k13_complex1 (k6_xcmplx_0 (k7_square_1 np_5) np_1) (k7_square_1 np_5))$$