

# l79\_sincos10 (TMHUXjJx- cXQ7vCnW6pj5MdfTqMxpFHWRXTJ)

October 27, 2020

Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k2\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_fdiff\_9 : \iota$  be given. Let  $k1\_rcomp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_real\_1 : \iota \Rightarrow \iota$  be given. Let  $k10\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k32\_sin\_cos : \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $np\_4 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_rcomp\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow ((r1\_tarski X0 (k9\_xtuple\_0 X1)) \Rightarrow (k9\_xtuple\_0 (k5\_relat\_1 X1 X0) = X0)) \quad (1)$$

Assume the following.

$$r1\_tarski (k3\_rcomp\_1 (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_2)) k6\_numbers) (k1\_relset\_1 k1\_numbers k2\_fdiff\_9) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1\_tarski X0 X1) \wedge (r1\_tarski X1 X2)) \Rightarrow (r1\_tarski X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((v1\_funct\_1 X2) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow (k2\_partfun1 X0 X1 X2 X3 = k5\_relat\_1 X2 X3) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (5)$$

Assume the following.

$$\begin{aligned} & r1\_tarski (k1\_rcomp\_1 (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_2)) \\ & (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_4))) (k3\_rcomp\_1 (k1\_real\_1 \\ & (k10\_real\_1 k32\_sin\_cos np\_2)) k6\_numbers) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1\_funct\_1 X2)\wedge \\ & (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))\Rightarrow((v1\_funct\_1 \\ & (k2\_partfun1 X0 X1 X2 X3))\wedge(m1\_subset\_1 (k2\_partfun1 X0 X1 X2 X3) \\ & (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (v1\_funct\_1 k2\_fdiff\_9)\wedge(m1\_subset\_1 k2\_fdiff\_9 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 k1\_numbers k1\_numbers))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \end{aligned} \quad (10)$$

**Theorem 1**

$$\begin{aligned} & k1\_relset\_1 k1\_numbers (k2\_partfun1 k1\_numbers k1\_numbers k2\_fdiff\_9 \\ & (k1\_rcomp\_1 (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_2)) (k1\_real\_1 \\ & (k10\_real\_1 k32\_sin\_cos np\_4)))) = k1\_rcomp\_1 (k1\_real\_1 (k10\_real\_1 \\ & k32\_sin\_cos np\_2)) (k1\_real\_1 (k10\_real\_1 k32\_sin\_cos np\_4)) \end{aligned}$$