

l80_enumset1
(TMaHXKmPk6Kjcyt2Ymsvr4mDCik83BEXsGa)

October 27, 2020

Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k2_enumset1\ X0\ X1\ X2\ X3 = k2_xboole_0\ (k1_enumset1\ X0\ X1\ X2)\ (k1_tarski\ X3) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k1_enumset1\ X0\ X1\ X2 = k1_enumset1\ X1\ X2\ X0 \quad (2)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\forall X3.k2_enumset1\ X0\ X1\ X2\ X3 = k2_enumset1\ X1\ X2\ X0\ X3$$