

180_finseq_3
(TMccVKGzCpwcsJKsqsdcedVcVEQmdzo8iia)

October 27, 2020

Let $k1_finseq_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(k1_finseq_3 X0 X1 = k1_xboole_0) \Leftrightarrow (r1_tarski (k10_xtuple_0 X0) X1)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow \\ & ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k3_finseq_1 X1 = np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (3)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. v1_finseq_1 (k5_finseq_1 X0) \quad (5)$$

Assume the following.

$$\forall X0. (v1_relat_1 (k5_finseq_1 X0)) \wedge (v1_funct_1 (k5_finseq_1 X0)) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. (k1_finseq_3 (k9_finseq_1 X0) X1 = k1_xboole_0) \Leftrightarrow \\ & (X0 \in X1) \end{aligned}$$