

## l80\_funct\_4

(TMV26eUdZgxPZEjYNfCkXY14RpBEuDbRhTW)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (r1\_partfun1 X0 X1))) \tag{1}$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_partfun1 X0 X1) \Leftrightarrow (k1\_funct\_4 X0 X1 = k1\_funct\_4 X1 X0))) \tag{2}$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_partfun1 X0 X1) \Leftrightarrow (k2\_xboole\_0 X0 X1 = k1\_funct\_4 X0 X1))) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Rightarrow (k2\_xboole\_0 X0 X1 = X1) \tag{4}$$

### Theorem 1

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (k1\_funct\_4 X1 X0 = X1)))$$