

186_fib_num4 (TM-
LaNRD2NoTLxPpDmqGTphh2E9UtCzPBaQV)

October 27, 2020

Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $np_5 : \iota$ be given. Let $np_3 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xreal_0 X0 X1) \Leftrightarrow (r1_xreal_0 (k4_xcmplx_0 X1) (k4_xcmplx_0 X0)))) \quad (1)$$

Assume the following.

$$((v2_xreal_0 np_5) \wedge (m2_subset_1 np_5 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_5 k5_numbers) \wedge (m1_subset_1 np_5 k1_numbers)) \quad (2)$$

Assume the following.

$$((v2_xreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \quad (3)$$

Assume the following.

$$\neg r1_xreal_0 np_3 (k7_square_1 np_5) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k7_square_1 X0) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1 $\neg r1_xreal_0 (k4_xcmplx_0 (k7_square_1 np_5)) (k4_xcmplx_0 np_3)$.