

# l8\_cfunct\_1

(TMMKE44LsdQthEj5N3cxZM6GemcrnHP9Fb)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $k8\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k8\_relset\_1 X0 X1 X2 X3 = k8\_relat\_1 X2 X3) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. (X2 = k8\_relat\_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in k9\_xtuple\_0 X0) \wedge (k1\_funct\_1 X0 X3 \in X1)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v5\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1))) \Rightarrow (\forall X2. (X2 \in k9\_xtuple\_0 X1) \Rightarrow (k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (6)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.(\neg v1\_xboole\_0 X2)\Rightarrow(\forall X3. \\ ((v1\_funct\_1 X3)\wedge(m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X2 \\ k2\_numbers))))\Rightarrow((X0 \in k8\_relset\_1 X2 k2\_numbers X3 X1)\Leftrightarrow((X0 \in k1\_relset\_1 \\ X2 X3)\wedge(k7\_partfun1 k2\_numbers X3 X0 \in X1))))$$