

l8_fib_fusc
(TMKhb6DnQAo8KMDEptaFeqtXLRJpSa8f81w)

October 27, 2020

Let $k10_ami_3 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg (X0 \neq X1) \wedge (k10_ami_3 X0 = k10_ami_3 X1))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (3)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (v7_ordinal1 X0) \quad (8)$$

Theorem 1 $k10_ami_3 k6_numbers \neq k10_ami_3 np_2$.