

l8_holder_1

(TMP5MpaFatZmdD1e9NbfCobY7CBS8yr4Vu5)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_series_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_valued_0 : \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_series_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v7_valued_0 X0) \Leftrightarrow (\forall X1. (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\ & (\forall X2. (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((r1_xxreal_0 \\ & X1 X2) \Rightarrow (r1_xxreal_0 (k8_nat_1 k1_numbers X0 X1) (k8_nat_1 k1_numbers \\ & X0 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\forall X1. (v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow ((v1_xboole_0 X0) \vee ((v2_xxreal_0 X1) \vee (v3_xxreal_0 X0)))))) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\forall X1. (v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v3_xxreal_0 X1)) \Rightarrow (v3_xxreal_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v2_xxreal_0 X0)) \Rightarrow (v2_xxreal_0 X1))) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ ((\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\ k6_numbers (k3_funct_2 k5_numbers k1_numbers X0 X1))) \Rightarrow (v7_valued_0 \\ (k3_series_1 X0))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(((v1_funct_1 X1) \wedge ((v1_funct_2 \\ X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ X0)))))) \wedge (v7_ordinal1 X2)) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2) \end{aligned} \quad (8)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (9)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (10)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ (k3_series_1 X0 = k2_series_1 X0) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge \\ (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1)))))) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_2 X0 \\ X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (13)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\ & k6_numbers (k1_seq_1 X0 X1))) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers \\ & k5_numbers) \Rightarrow (r1_xxreal_0 (k1_seq_1 X0 X1) (k1_seq_1 (k3_series_1 \\ & X0) X1)))) \end{aligned} \quad (15)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (v1_xreal_0 (k1_funct_1 X0 X1)) \quad (17)$$

Assume the following.

$$v3_membered k1_numbers \quad (18)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (19)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (20)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (21)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v1_funct_1 (k3_series_1 X0)) \wedge ((v1_funct_2 (k3_series_1 X0) \\ & k5_numbers k1_numbers) \wedge (m1_subset_1 (k3_series_1 X0) (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers k1_numbers)))))) \end{aligned} \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (23)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v3_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xreal_0 X0) \wedge (\neg v2_xxreal_0 X0))) \quad (24)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v2_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X0))) \quad (25)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (26)$$

Assume the following.

$$\forall X0. \forall X1. (v3_membered X1) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v3_valued_0 X2)) \quad (27)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((\forall X1. (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (r1_xxreal_0 \\ & k6_numbers (k1_seq_1 X0 X1))) \Rightarrow (\forall X1. (m2_subset_1 X1 k1_numbers \\ & k5_numbers) \Rightarrow ((k1_seq_1 (k3_series_1 X0) X1 = k6_numbers) \Rightarrow (\forall X2. \\ & (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((r1_xxreal_0 X2 X1) \Rightarrow \\ & (k1_seq_1 X0 X2 = k6_numbers)))))) \end{aligned}$$