

l8_series_1 (TMUHAcWEKGF- puS7gu8TG8niXev5HdVKr2m7)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k1_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X0 = X0 \quad (6)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (7)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (8)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (v1_partfun1 X1 X0) \Leftrightarrow (k1_relset_1 X0 X1 = X0) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 X1))) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X2 = k1_valued_1 X0 X1) \Leftrightarrow ((k9_xtuple_0 X2 = k3_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \wedge (\forall X3. (X3 \in k9_xtuple_0 X2) \Rightarrow (k1_funct_1 X2 X3 = k2_xcmplx_0 (k1_funct_1 X0 X3) (k1_funct_1 X1 X3))))))) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_partfun1 X0 k5_numbers) \wedge (v1_valued_0 X0)))))) \Rightarrow (\forall X1. \\ & ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 k5_numbers) \wedge ((v1_funct_1 X1) \wedge ((v1_partfun1 X1 k5_numbers) \wedge (v1_valued_0 X1)))))) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge ((v4_relat_1 X2 k5_numbers) \wedge ((v1_funct_1 X2) \wedge ((v1_partfun1 X2 k5_numbers) \wedge (v1_valued_0 X2)))))) \Rightarrow ((X0 = k1_valued_1 X1 X2) \Leftrightarrow (\forall X3. (m2_subset_1 X3 k1_numbers k5_numbers) \Rightarrow (k1_funct_1 X0 X3 = k2_xcmplx_0 (k1_funct_1 X1 X3) (k1_funct_1 X2 X3)))))) \end{aligned}$$