

l90_xxreal_3

(TMWyd7RpqrWVMBE8t6bEA4SYtQYF91JLCgp)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k4_xxreal_3 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k4_xxreal_3 X0 k6_numbers = k6_numbers) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v1_xxreal_0 (k4_xxreal_3 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k4_xxreal_3 X0 X1 = k4_xxreal_3 X1 X0) \quad (3)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((X0 = k6_numbers) \Rightarrow (k4_xxreal_3 X0 (k4_xxreal_3 X1 X2) = k4_xxreal_3 (k4_xxreal_3 X0 X1) X2))))$$