

193_funct_7 (TMTHKEp-
gUnPQ7mvFXEJLEhw4XJdW2Wc6mJT)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k5_numbers) \Rightarrow (\forall X2.(v1_relat_1 X2) \Rightarrow (k3_relat_1 (k9_funct_7 \\ & X2 X0) (k9_funct_7 X2 X1) = k9_funct_7 X2 (k2_nat_1 X1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k9_funct_7 \\ & X0 (k1_nat_1 X1 np_1) = k3_relat_1 X0 (k9_funct_7 X0 X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_xcmplx_0 np_1 X0 = X0) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1_xcmplx_0 X0) \wedge ((v1_xcmplx_0 \\ & X1) \wedge (v1_xcmplx_0 X2))) \Rightarrow (k3_xcmplx_0 (k2_xcmplx_0 X0 X1) X2 = k2_xcmplx_0 \\ & (k3_xcmplx_0 X0 X2) (k3_xcmplx_0 X1 X2)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k4_nat_1 X0 X1 = k3_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_subset_1 X1 k5_numbers))\Rightarrow(k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (10)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v1_relat_1 (k9_funct_7 X0 X1)) \quad (12)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(m2_subset_1 (k4_nat_1 X0 X1) k1_numbers k5_numbers) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(m2_subset_1 (k2_nat_1 X0 X1) k1_numbers k5_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k4_nat_1 X0 X1 = k4_nat_1 X1 X0) \quad (16)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (18)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (20)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k5_numbers) \Rightarrow (\forall X2.(v1_relat_1 X2) \Rightarrow ((k9_funct_7 (k9_funct_7 \\ & X2 X0) X1 = k9_funct_7 X2 (k4_nat_1 X0 X1)) \Rightarrow (k9_funct_7 (k9_funct_7 \\ & X2 X0) (k2_nat_1 X1 np_1) = k9_funct_7 X2 (k4_nat_1 X0 (k2_nat_1 \\ & X1 np_1)))))) \end{aligned}$$