

l95\_toprealb  
(TMVzyNsytwfrqGVLo83yPydvxqxYVVtnewC)

October 27, 2020

Let  $k1\_pre\_topc : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_topalg\_2 : \iota$  be given. Let  $k2\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $k3\_topmetr : \iota$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. ((v1\_pre\_topc X0) \wedge (l1\_pre\_topc X0)) \Rightarrow (k1\_pre\_topc X0 (k2\_struct\_0 X0) = X0) \quad (1)$$

Assume the following.

$$k2\_topalg\_2 = k3\_topmetr \quad (2)$$

Assume the following.

$$(\neg v2\_struct\_0 k3\_topmetr) \wedge ((v1\_pre\_topc k3\_topmetr) \wedge (v2\_pre\_topc k3\_topmetr)) \quad (3)$$

Assume the following.

$$(v2\_pre\_topc k3\_topmetr) \wedge (l1\_pre\_topc k3\_topmetr) \quad (4)$$

**Theorem 1**  $k1\_pre\_topc k2\_topalg\_2 (k2\_struct\_0 k2\_topalg\_2) = k2\_topalg\_2.$