

197_jordan
(TMNPPmvy9SVjtzHhDNCitPSHFmsrzteikjH)

October 27, 2020

Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_3 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1 $k17_euclid (k19_euclid k6_numbers np_3) = k6_numbers$.